

Dynamics of broadband dispersive Alfvén waves

M. Marklund and P.K. Shukla

Department of Physics, Umeå University, SE-901 87 Umeå, Sweden

(Dated: February 2, 2008)

Abstract

The properties of amplitude modulated broadband Alfvén waves is investigated. In particular, the dynamics of circularly polarized dispersive Alfvén waves, governed by a derivative nonlinear Schrödinger equation, is analyzed using the Wigner formalism. The modulational instability of random phase dispersive pump Alfvén waves is investigated, and it is shown that the spectral broadening gives rise to a new mode structure.

PACS numbers: 52.35.Mw, 52.40.Db, 94.30.Tz

I. INTRODUCTION

About thirty years ago, Rogister [1] introduced an elegant perturbation technique of the Vlasov–Maxwell system of equations in order to investigate amplitude modulation of magnetic field-aligned circularly polarized dispersive Alfvén waves in an electron–ion plasma. He derived a derivative nonlinear Schrödinger (DNLS) equation, which is exactly integrable by the inverse scattering transform method [2]. The DNLS equation for nonlinear dispersive Alfvén waves have also been derived by means of the two-fluid – Maxwell system of equations [3, 4, 5, 6, 7, 8], and the modulational instability of a constant amplitude Alfvén pump wave has been investigated. Also reported is the algebraic envelope Alfvén soliton [3, 9].

The DNLS equation has been extended to include the effects of collisions and wave-particle interactions [10, 11]. The resulting modified DNLS equation has been used to investigate damping of the envelope Alfvénic soliton.

In this paper, we report on the nonlinear stability of broadband dispersive Alfvén waves in plasmas. We introduce a Wigner–Moyal formalism [12, 13] for the modified DNLS equation and derive a kinetic equation for Alfvénic quasiparticles. While the DNLS equation is appropriate for studies of modulational instability of a coherent Alfvén pump wave, the kinetic equation can be adopted to investigate the effects of partial coherence and random phases associated with broadband dispersive Alfvén wave packets.

II. GOVERNING EQUATIONS

Let us consider the nonlinear propagation of circularly polarized dispersive Alfvén waves along the external magnetic field $B_0\hat{\mathbf{z}}$, where B_0 is the magnetic field strength and $\hat{\mathbf{z}}$ is the unit vector along the z -axis. The dynamics of modulated Alfvén wave envelopes is governed by the modified DNLS equation [10, 11]

$$i\partial_t u + \partial_z^2 u + is\partial_z(|u|^2 u) = p[u, u^*], \quad (1)$$

where the subscripts t and z denotes the corresponding derivatives. Equation (1) governs the amplitude modulation of circularly polarized Alfvén waves. Here $u = (B_x + iB_y)/(2|1 - \beta|^{1/2}B_0)$, $\beta = V_s^2/V_A^2$, $s = \text{sign}(1 - \beta)$, time and space coordinates are normalized by the ion gyroperiod ω_{ci}^{-1} and the ion skin depth $V_A/\omega_{ci} = c/\omega_{pi}$. For our purposes, we have in

the collision-dominated case

$$p = iD\partial_z^2 u, \quad (2)$$

where $D = (1/2)[(\eta_1/\rho_0) + c^2\eta_2/4\pi]\omega_{ci}/V_A^2$, η_1 is the coefficient of the ion viscosity and η_2 is the resistivity. On the other hand, in a collisionless case, we have

$$p = i\frac{C}{4\pi}\partial_z(uH(z, t)) = i\frac{C}{4\pi}\partial_z\left(u\text{P}\int_{-\infty}^{+\infty} dz' \frac{|u(z', t)|^2}{z' - z}\right), \quad (3)$$

where P denotes the principal value. The coefficient C depends on the velocity distribution of the particle species. For $V_A \gg V_s$, we have

$$C = \sqrt{\frac{m_e}{2\pi m_i}} \left(\frac{V_s}{V_A}\right) \exp\left(-\frac{V_A^2}{2V_{Te}^2}\right) \ll 1, \quad (4)$$

where V_{Te} is the electron thermal speed.

With (2) and (3), Eq. (1) is a special case of the equation

$$i\partial_t u + g\partial_z^2 u + F(|u|^2; t, z)u + i\partial_z[G(|u|^2; t, z)u] = 0, \quad (5)$$

where g is a complex valued constant and F and G are complex valued functions. This equation can be found in a wide variety of applications, such as Bose-Einstein condensation and nonlinear optics, apart from in plasma physics. Next we proceed to analyze Eq. (5) for broad band waves.

In order to study the stability of broad band waves governed by Eq. (5), it is convenient to introduce the Wigner function

$$\rho(t, z, k) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\zeta e^{ik\zeta} \langle u^*(t, z + \zeta/2) u(t, z - \zeta/2) \rangle \quad (6)$$

for the Alfvén waves. Here the angular brackets represents the ensemble average. The Wigner function is given by the Fourier transform of the two-point correlation function, and as a generalized distribution function for quasi-particles it can be used to describe a broad band spectrum of the field represented by u , e.g. Alfvén waves. We note that from (6) the equality

$$I(t, z) \equiv \langle |u(t, z)|^2 \rangle = \int_{-\infty}^{\infty} dk \rho(t, z, k) \quad (7)$$

holds. Applying the time derivative to the Wigner function (6) and using (5), one obtains the kinetic equation

$$\begin{aligned} \partial_t \rho + 2\text{Re}(g)k\partial_z \rho + \text{Im}(g)\left(\frac{1}{2}\partial_z^2 - 2k^2\right)\rho + 2\text{Im}\left[F \exp\left(\frac{i}{2}\overleftarrow{\partial}_z \overrightarrow{\partial}_k\right)\right]\rho \\ + \partial_z \left\{ \text{Re}\left[G \exp\left(\frac{i}{2}\overleftarrow{\partial}_z \overrightarrow{\partial}_k\right)\right]\rho \right\} - 2k\text{Im}\left[G \exp\left(\frac{i}{2}\overleftarrow{\partial}_z \overrightarrow{\partial}_k\right)\right]\rho = 0. \end{aligned} \quad (8)$$

As an example of the versatility of Eq. (8), we look at the case $g = 1$, $F = I$, and $G = 0$. Then Eq. (5) reduces to the regular NLSE, and Eq. (8) becomes

$$\partial_t \rho + 2k \partial_z \rho + 2I \sin \left(\frac{i}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \rho = 0, \quad (9)$$

an equation which to lowest order gives the Vlasov equation

$$\partial_t \rho + 2k \partial_z \rho + (\partial_z I)(\partial_k \rho) = 0. \quad (10)$$

For Eq. (1) we have $F = 0$, $g = 1 - iD$, and $G = sI - (C/4\pi)H$, and the kinetic equation (8) takes the form

$$\partial_t \rho + 2k \partial_z \rho - 2skI \sin \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \rho + s \partial_z \left[I \cos \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \rho \right] = \hat{L} \rho, \quad (11)$$

where the operator expression on the right hand side is defined according to

$$\hat{L} \rho = D \left(\frac{1}{2} \partial_z^2 - 2k^2 \right) \rho \quad (12a)$$

in the collisional case (2), and

$$\begin{aligned} \hat{L} \rho = & -\frac{C}{2\pi} k \left[\text{Im}(H) \cos \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) + \text{Re}(H) \sin \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \right] \rho \\ & + \frac{C}{4\pi} \partial_z \left[-\text{Im}(H) \sin \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \rho + \text{Re}(H) \cos \left(\frac{1}{2} \overleftarrow{\partial}_z \overrightarrow{\partial}_k \right) \rho \right] \end{aligned} \quad (12b)$$

in the collisionless case (3).

III. ANALYSIS OF THE COHERENT CASE

We note that Eq. (1) has solutions of the modified plane wave form $u(t, z) = \bar{u}(t) \exp[ik_0 z - i\omega_0(t)t]$. In the collisional case (2) the time-dependent frequency is given by

$$\omega_0(t) = k_0^2 + \frac{su_0^2}{2Dk_0 t} [1 - \exp(-2Dk_0^2 t)], \quad (13a)$$

while the amplitude is exponentially damped according to

$$\bar{u}(t) = u_0 \exp(-Dk_0^2 t). \quad (13b)$$

Here u_0 is the constant amplitude of the solution, and we note that as $D \rightarrow 0$ we obtain the time-independent dispersion relation $\omega_0 = k_0^2 + sk_0 u_0^2$. In the collisionless case (3) however,

the character of the function $p[u, u^*]$ together with the above plane wave ansatz makes the amplitude time-independent, and subsequently the frequency becomes

$$\omega_0 = k_0^2 + sk_0 u_0^2, \quad (14)$$

where $\bar{u} = u_0$ denotes the constant amplitude. Thus, with the plane wave ansatz, the effects due to the thermal correction (3) vanish.

Coherent modulational instabilities can be analyzed by letting $u = (\bar{u}(t) + u_1(t, z)) \exp[ik_0 z - i\omega_0(t)t] + \text{c.c.}$, where $|u_1| \ll \bar{u}$, the time variation of $\bar{u}$ is assumed slow compared to the time scale of the perturbation, and c.c. denotes the complex conjugate. Linearizing Eq. (1) with respect to u_1 , dividing u_1 into its real and imaginary part, and assuming the wavenumber and frequency K and Ω of the perturbation we find the dispersion relation

$$\Omega = -iDK^2 + 2k_0K + 2sK\bar{u}^2 \pm [s^2K^2\bar{u}^4 + K^4 + 2sk_0K^2\bar{u}^2 - 4Dk_0K(Dk_0K + iK^2 + isk_0\bar{u}^2)]^{1/2} \quad (15a)$$

in the collisional case (2), and

$$\Omega = -\frac{1}{4}iCK\bar{u}^2 + 2k_0K + 2sK\bar{u}^2 \pm [s^2K^2\bar{u}^4 + K^4 + 2sk_0K^2\bar{u}^2 - \frac{1}{2}CK^2\bar{u}^2 (\frac{1}{8}C\bar{u}^2 + ik_0 + is\bar{u}^2)]^{1/2} \quad (15b)$$

in the collisionless case (3). We note from (15a) and (15b) that for $s = 1$ there is no instability and the perturbation modes are always damped. However, for $s = -1$ this is not the case. Letting $\gamma = \text{Im}(\Omega)$ where γ is the growth rate, we find from (15a)

$$\gamma = -DK^2 + (2k_0K^2\bar{u}^2 - K^2\bar{u}^4 - K^4)^{1/2}, \quad (16)$$

if we assume that $D \ll 1$ and linearize for D . Similarly, in the collisionless case, the parameter $C \ll 1$, and we may expand (15b), keeping terms linear in C . Then we find

$$\gamma = -\frac{1}{4}CK\bar{u}^2 + (2k_0K^2\bar{u}^2 - K^2\bar{u}^4 - K^4)^{1/2}. \quad (17)$$

Thus, the expressions (16) and (17) show that for $s = -1 \Leftrightarrow V_s > V_A$ we have growing modes due to the nonlinear evolution of the Alfvén waves. However, these modes suffer damping due to dissipative processes.

Next we will compare the above results for the coherent modulational instability with the case of partial coherent Alfvén waves.

IV. EFFECTS OF PARTIAL COHERENCE

Returning to Eq. (11), we first look for solutions of the form $\rho = \bar{\rho}(t, k)$. In the collisional case (12a) Eq. (11) takes the form $\partial_t \bar{\rho} + 2Dk^2 \bar{\rho} = 0$. This is integrated to yield

$$\bar{\rho} = \rho_0(k) \exp(-2Dk^2 t). \quad (18)$$

Equation (18) is in agreement with the monochromatic result (13b) based on Eq. (1), if we let $\rho_0(k) = I_0 \delta(k - k_0)$. In the collisionless case (3), the ansatz for ρ gives $H = 0$, where $\bar{I}(t) = \int dk \bar{\rho}(t, k)$. Thus, the expression (12b) gives $\bar{\rho} = \rho_0(k)$ through Eq. (11), in agreement with the monochromatic result.

Next we look at perturbation around the background solutions. Thus, we let $\rho = \bar{\rho}(t, k) + \rho_1(k) \exp(iKz - i\Omega t) + \text{c.c.}$, where $|\rho_1| \ll \rho_0$ and $|\partial_t \bar{\rho}| \ll \Omega \bar{\rho}$, and linearize with respect to ρ_1 . With $I = \bar{I}(t) + I_1 \exp(iKz - i\Omega t) + \text{c.c.}$ it follows that $H \approx \bar{H} + H_1 = H_1 = i\pi I_1 \exp(iKz - i\Omega t) + \text{c.c.}$ Thus, using

$$2 \sin\left(\frac{i}{2} K \partial_k\right) \bar{\rho}(t, k) = i [\bar{\rho}(t, k + K/2) - \bar{\rho}(t, k - K/2)] \quad (19a)$$

and

$$2 \cos\left(\frac{i}{2} K \partial_k\right) \bar{\rho}(t, k) = \bar{\rho}(t, k + K/2) + \bar{\rho}(t, k - K/2), \quad (19b)$$

we find that the nonlinear dispersion relation

$$1 = \frac{(s - ia)}{2K} \int dk \frac{(k - K/2) \bar{\rho}(t, k + K/2) - (k + K/2) \bar{\rho}(t, k - K/2)}{k + [sK\bar{I}(t) - \Omega - iL(k)]/2K} \quad (20)$$

valid for partially coherent Alfvén waves. Here $L(k) = D(K^2/2 + 2k^2)$ and $a = 0$ in the collisional case (2), and $L(k) = 0$ and $a = C/4$ in the collisionless case (3). In the monochromatic case, we have $\rho_0(k) = I_0 \delta(k - k_0)$. The dispersion relation (20) then becomes

$$\Omega = -\frac{1}{2}i(L_+ + L_-) + 2k_0 K + (2s - ia)K\bar{I} \quad (21)$$

$$\pm \left\{ (s - ia)^2 K^2 \bar{I}^2 + K^4 + 2(s - ia)k_0 K^2 \bar{I} - (L_+ - L_-) \left[\frac{1}{4}(L_+ - L_-) + iK^2 + i(s - ia)k_0 \bar{I} \right] \right\}^{1/2}$$

where $L_{\pm} = L(k_0 \pm K/2)$. Thus, inserting the definitions of $L(k)$ and a we regain the dispersion relations (15) when neglecting the slow background time variations.

However, if the Alfvén waves suffer random perturbations in e.g. the phase, the correlation between the waves may be nontrivial [14]. The partial coherence introduced by a random phase may be modeled by the Lorentzian distribution [15]

$$\rho_0(k) = \frac{I_0}{\pi} \frac{\Delta}{(k - k_0)^2 + \Delta^2}, \quad (22)$$

where Δ denotes the width of the distribution around k_0 . Due to the finite width of the Lorentz distribution the dispersion integral (20) will have poles. In the collisional case (2) we obtain a rather lengthy analytical expression, where a complicated interplay between the spectral broadening, represented by Δ , and the collision parameter D takes place. Thus, we do not explicitly state this particular result here, but instead give the numerical solution of (20) in conjunction with the Lorentzian distribution (22) in the collisional case (2) (see Fig. 1). However, in the collisionless case (3), the result is somewhat more compact and takes the form

$$1 = 2KI_0 \left(s - \frac{1}{4}iC \right) \left\{ \frac{\bar{\Omega} + k_0K + iK\Delta}{(\bar{\Omega} + 2iK\Delta)^2 - K^4} - iK\Delta \frac{4K^2\Delta^2 + (\bar{\Omega} - K^2)^2/2 + (\bar{\Omega} + K^2)^2/2 + 2(\bar{\Omega} + 2k_0K)(\bar{\Omega} - sKI_0)}{[4K^2\Delta^2 + (\bar{\Omega} - K^2)^2][4K^2\Delta^2 + (\bar{\Omega} + K^2)^2]} \right\}, \quad (23)$$

where $\bar{\Omega} = \Omega - 2k_0K - sKI_0$.

Letting $\bar{\Omega} = \text{Re}(\bar{\Omega}) + i\Gamma$, where Re denotes the real part, we can solve the nonlinear dispersion relations numerically for the growth rate Γ as a function of the wavenumber K . The results are shown in Figs. 1 and 2, where we have used $I_0 = 0.5$ and $k_0 = 1$. We see that the parameter region with $s = 1$ exhibits new properties as compared to the coherent case. A new type of instability occurs due to the spectral broadening of the dispersive Alfvén pump wave.

V. SUMMARY

We have investigated the nonlinear stability of broadband dispersive Alfvén waves in magnetoplasmas, using a Wigner–Moyal formalism. A new mode structure with new instabilities, due to the finite spectral width of the dispersive Alfvén pump wave, is found. The spectral broadening in conjunction with the kinetic modification of the DNLS equation can thus give rise to growing modes not present in the coherent case.




FIG. 1: The growth rate for different parameter values in the collisional case (2). The upper left panel has $s = -1$, while $D = \Delta = 0$; the middle upper panel has $s = -1$ and $D = 0$, while $\Delta = 0.1$; the right upper panel has $s = -1$ and $\Delta = 0$, while $D = 0.1$. The left lower panel has $s = -1$ and $D = \Delta = 0.1$, while the right panel has $s = 1$ and $D = \Delta = 0.1$. Note that the last of these panels has no coherent counterpart.




FIG. 2: The growth rate for different parameter values in the collisional case (3). The upper left panel has $s = -1$, while $C = \Delta = 0$; the middle upper panel has $s = -1$ and $C = 0$, while $\Delta = 0.1$; the right upper panel has $s = -1$ and $\Delta = 0$, while $C = 0.1$. The left lower panel has $s = -1$ and $C = \Delta = 0.1$, while the right panel has $s = 1$ and $C = \Delta = 0.1$. Note that the last of these panels has no coherent counterpart.

-
- [1] A. Rogister, Phys. Fluids **14**, 2733 (1971).
 - [2] D. J. Kaup and A. C. Newell, J. Math. Phys. **19**, 798 (1978).
 - [3] E. Mjølhus, J. Plasma Phys. **16**, 321 (1976).
 - [4] V. M. Lashkin, Phys. Rev. E **69**, 016611 (2004).
 - [5] K. Mio, T. Ogino, K. Minami, and S. Takeda, J. Phys. Soc. Jpn. **41**, 265 (1976).
 - [6] S. R. Spangler and J. P. Sheerin, J. Plasma Phys. **27**, 193 (1982).
 - [7] J.-I. Sakai and B. U. O. Sonnerup, J. Geophys. Res. **88**, 9069 (1983).
 - [8] M. S. Ruderman, Phys. Plasmas **9**, 2940 (2002).
 - [9] T. Hada, C. F. Kennel, and B. Buti, J. Geophys. Res. **94**, 65 (1989).
 - [10] M. V. Medvedev and P. H. Diamond, Phys. Plasmas **3**, 863 (1996).
 - [11] M. V. Medvedev, V. I. Shevchenko, P. H. Diamond, and V. L. Galinsky, Phys. Plasmas **4**, 1257 (1997).
 - [12] E. P. Wigner, Phys. Rev. **40**, 749 (1932).
 - [13] J. E. Moyal, Proc. Cambridge Philos. Soc. **45**, 99 (1949).
 - [14] Yu. L. Klimontovich, *The Statistical Theory of Non-Equilibrium Processes in a Plasma* (Pergamon Press, Oxford, 1967).
 - [15] R. Loudon, *The Quantum Theory of Light* (Oxford University Press, Oxford, 2000).